

REGULAR LIE GROUPS AND A THEOREM OF LIE-PALAIS

VLADIMIR G. PESTOV

Victoria University of Wellington,
P.O. Box 600, Wellington, New Zealand

vladimir.pestov@vuw.ac.nz

ABSTRACT. In 1984 Milnor had shown how to deduce the Lie-Palais theorem on integration of infinitesimal actions of finite-dimensional Lie algebras on compact manifolds from general theory of regular Lie groups modelled on locally convex spaces. We show how, in the case of effective action, one can eliminate from Milnor's argument the abstract Lie-Cartan theorem, making the deduction rather elementary. A machinery employed in the proof provides a partial solution to a problem examined in 1972 by van Est and Świerczkowski.

1. Introduction. A well-known result dated back to Lie and finalized by Palais [Pa] states that every infinitesimal action of a finite-dimensional Lie algebra on a compact smooth manifold, X , is derived from a smooth action of a finite-dimensional Lie group on X . Milnor [M] gives a proof of this theorem based on theory of regular Lie groups. This proof, however, is only partial (in Milnor's own words), not being self-contained: by necessity, it invokes the abstract Lie-Cartan theorem. One definitely cannot hope to circumvent this landmark result while proving the Lie-Palais theorem (which simply turns into the Lie-Cartan theorem in the degenerate case of a trivial action). However, things are different in the most important particular case where the infinitesimal action is effective, that is, one deals with finite-dimensional Lie algebras of vector fields. We show how under this assumption the Lie-Cartan theorem can be eliminated from the Milnor's argument, which therefore becomes quite elementary in the sense that it does not invoke anything advanced beyond the scope of theory of regular Lie groups (like the structure theory of Lie algebras).

Call a Banach-Lie algebra coming from a Lie group *enlargable*. The Lie-Cartan theorem essentially says that every finite-dimensional Lie algebra is enlargable; in infinite dimensions non-enlargable Banach-Lie algebras can be found [vEK]. However, a Banach-Lie algebra is enlargable as soon as it admits a continuous monomorphism into an enlargable Banach-Lie algebra [H, Ś1, vEK].

For more general Lie groups and algebras the situation is much harder to deal with. A closed Lie subalgebra of the Lie algebra of a Fréchet-Lie group G need

1991 *Mathematics Subject Classification.* Primary 22E65; Secondary 17B66, 58D05.

Key words and phrases. Enlargable Banach-Lie algebras, regular Lie groups, integrability, Lie algebras of vector fields, Lie-Palais theorem.

not have an associated Fréchet-Lie group even if G meets certain conditions of regularity [KYMO, M]. For some positive advances in this direction, see [L].

We show that the above result on Banach-Lie algebras can be pushed somewhat further to absorb a restricted version of the Lie-Palais theorem. Let \mathfrak{h} be the Lie algebra of a regular Lie group modelled on a locally convex space, and let \mathfrak{g} be a Banach-Lie algebra with finite-dimensional center admitting a continuous monomorphism into \mathfrak{h} . Then \mathfrak{g} is enlargable. The result is deduced from regular Lie group theory by means of the concept of a free Banach-Lie algebra introduced by us earlier [Pe1]. In passing, we observe that this concept provides a partial answer to a problem investigated in 1972 by van Est and Świerczkowski [vEŚ].

Since the group of diffeomorphisms of a compact manifold forms a regular Fréchet-Lie group, an application of our main result yields a theorem on integration of finite-dimensional Lie algebras of vector fields. The latter theorem thus merges fully in the realm of regular Lie group theory.

2. Preliminaries. We first recall a few basic facts about regular Lie groups. A C^∞ (Gâteaux) smooth Lie group G modelled on a bornological sequentially complete locally convex space is called *regular* if every smooth path $v : I = [0, 1] \rightarrow \text{Lie}(G)$ has a left product integral $p : I \rightarrow G$, that is, a solution to the differential equation $Dp(t) = p(t) \cdot v(t)$, and if furthermore the correspondence $v \mapsto p(0)^{-1} \cdot p(1)$ defines a smooth map from the locally convex space $C^\infty(I, \text{Lie}(G))$ to G . (This definition is taken from Milnor's survey [M], where regularity is understood in a somewhat less restrictive sense than by Kobayashi *et al* [KYMO]. Notation is self-explanatory.)

The condition of regularity seems indispensable if one wants to derive substantial results, and all known examples of Lie groups modelled on locally convex spaces are regular. Such are: the groups of diffeomorphisms of smooth compact manifolds, certain subgroups of these, the groups of currents (and their central extensions), Banach-Lie groups, and many more [KYMO, M]. The following is a basic result about regular Lie groups.

Theorem 2.1 ([M]; cf. [KYMO]). *Let G and H be smooth Lie groups modelled on bornological sequentially complete locally convex spaces. Let G be simply connected and H be regular. Then every continuous Lie algebra morphism $\text{Lie}(G) \rightarrow \text{Lie}(H)$ is tangent to a (necessarily unique) Lie group morphism $G \rightarrow H$. \square*

We also need a test for enlargability of Banach-Lie algebras. Recall that every such algebra carries a natural structure of a group germ [Ś3].

Theorem 2.2 [H, Ś1]. *Let \mathfrak{g} be a Banach-Lie algebra such that there exist a group G and a map $\phi : \mathfrak{g} \rightarrow G$ which is locally one-to-one and a morphism of group germs. Then \mathfrak{g} is enlargable. Moreover, G can be given a unique structure of a Banach-Lie group associated to \mathfrak{g} in such a way that ϕ becomes an exponential map. \square*

3. Free Banach-Lie algebras.

Theorem 3.1 [Pe1]. *Let E be a normed space. There exist a complete normed Lie algebra $\mathcal{FL}(E)$ and a linear isometry $i_E : E \hookrightarrow \mathcal{FL}(E)$ such that:*

1. *The image of i_E topologically generates $\mathcal{FL}(E)$.*
2. *For every complete normed Lie algebra \mathcal{L} and an arbitrary contracting linear operator $f : E \rightarrow \mathcal{L}$ there exists a contracting Lie algebra homomorphism $\hat{f} : \mathcal{FL}(E) \rightarrow \mathcal{L}$ with $\hat{f} \circ i_E = f$.*

The pair $(\mathcal{FL}(E), i_E)$ with the properties 1 and 2 is unique up to an isometrical isomorphism. \square

We call $\mathcal{FL}(E)$ the *free Banach-Lie algebra* on E . It is easy to prove that if $\dim E \geq 2$ then $\mathcal{FL}(E)$ is centerless.

Let Γ be a set. We denote the Banach-Lie algebra $\mathcal{FL}(l_1(\Gamma))$ simply by $\mathcal{FL}(\Gamma)$ and call the free Banach-Lie algebra on a set Γ . It is easy to see that the above algebra possesses the following universal property: every map f sending Γ to the unit ball of a Banach-Lie algebra \mathfrak{g} extends in a unique way to a contracting Lie algebra morphism from $\mathcal{FL}(\Gamma)$ to \mathfrak{g} . The well-known fact that a Banach space of density character $\leq \text{Card}(\Gamma)$ is a factor space of $l_1(\Gamma)$ implies that an arbitrary Banach-Lie algebra is a factor Banach-Lie algebra of a free Banach-Lie algebra of the form $\mathcal{FL}(\Gamma)$ [Pe1].

Remark 3.2. Not every Banach-Lie algebra \mathfrak{g} admits a faithful representation in a Banach space, even if \mathfrak{g} is enlargable [vEŠ]. An investigation [vEŠ] of van Est and Świerczkowski was partly motivated by the problem on whether or not every Banach-Lie algebra is a quotient algebra of a Banach-Lie algebra with such a property. It was suggested that the algebra $\Lambda \mathfrak{g}$ of continuous paths starting at zero (which is enlargable and of which \mathfrak{g} is a quotient) can be represented in $\mathfrak{gl}(E)$ for some E . While this conjecture remains unproved, the following result can be considered as one possible answer to the above motivating question. (This fact was left unnoticed by us earlier in [Pe1].)

Theorem 3.3. *Every Banach-Lie algebra is a factor algebra of a Banach-Lie algebra admitting a faithful representation in a Banach space.*

Proof. The left regular representation of a free Banach-Lie algebra $\mathcal{FL}(E)$ is faithful if $\dim E \geq 2$. \square

Theorem 3.4 [Pe1]. *Every free Banach-Lie algebra on a normed space is enlargable.*

Proof. A Banach-Lie algebra admitting a faithful representation in a Banach space is enlargable [vEK]. \square

Corollary 3.5 [Š2]. *Every Banach-Lie algebra is a factor algebra of an enlargable Banach-Lie algebra.* \square

We shall denote by $\mathcal{FG}(\Gamma)$ the simply connected Banach-Lie group attached to the free Banach-Lie algebra $\mathcal{FL}(\Gamma)$. It is couniversal among all connected Banach-Lie groups of density character $\leq \text{Card}(\Gamma)$ [Pe1].

4. Extension of Lie-Palais theorem in case of effective action.

Lemma 4.1. *Let \mathfrak{g} be a Banach-Lie algebra with finite-dimensional center \mathfrak{z} . Let f be a continuous map from \mathfrak{g} to a Hausdorff topological group G which is a morphism of group germs. Suppose that for every $x \in \mathfrak{z}$ there is a $\lambda \in \mathbb{R}$ with $f(\lambda x) \neq e_G$. Then \mathfrak{g} is enlargable.*

Proof. The set $\mathfrak{z} \cap \ker f$ forms a closed additive subgroup of a finite-dimensional vector space, containing no one-dimensional linear subspaces; in other words, it is a discrete lattice, and there exists an open neighbourhood of zero, U , in \mathfrak{z} such that $U \cap \ker f = \{0\}$. Choose an open $\tilde{U} \subset \mathfrak{g}$ with $0 \in \tilde{U} \cap \mathfrak{z} \subset U$.

Let $\phi: \mathfrak{g} \rightarrow F$ be a universal morphism of a group germ to a group. (That is, any other such morphism, $g: \mathfrak{g} \rightarrow A$, is a composition of ϕ and a group homomorphism $F \rightarrow A$; cf. [Š3].) Let $x \in \tilde{U} \setminus \{0\}$; it suffices to prove that $\phi(x) \neq e_G$ and apply Theorem 2.2. Now, if $x \notin \mathfrak{z}$, then x is separated from zero by the composition of the adjoint representation $x \mapsto \text{ad}_x$ and the exponentiation $\text{End}(\mathfrak{g}_+) \rightarrow \text{GL}(\mathfrak{g}_+)$ (both being morphisms of group germs). If $x \in \mathfrak{z}$, then $x \in U$ and $f(x) \neq e_G$. \square

Main Theorem 4.2. *Let \mathfrak{g} be a Banach-Lie algebra with finite-dimensional center and let \mathfrak{g} admit a continuous monomorphism into the Lie algebra of a regular Lie group. Then \mathfrak{g} is enlargable.*

Proof. Let H be a regular Lie group and $h: \mathfrak{g} \rightarrow \text{Lie}(H)$ be a continuous Lie monomorphism. Denote by B the unit ball in \mathfrak{g} , and by i the contracting Lie algebra morphism $\mathcal{FL}(B) \rightarrow \mathfrak{g}$ extending the identity map $B \rightarrow B$. According to Theorem 2.1, the composition morphism $h \circ i: \mathcal{FL}(B) \rightarrow \text{Lie}(H)$ is tangent to a morphism of Lie groups $\hat{i}: \mathcal{FG}(B) \rightarrow H$. Denote $I = \ker h \circ i$, $J = \ker \hat{i}$, and let A be a subgroup of $\mathcal{FG}(B)$ algebraically generated by the exponential image of I . Let G denote the Hausdorff topological group $\mathcal{FG}(B)/J$, and let $f: \mathfrak{g} \rightarrow G$ be a map defined by $f(x + I) = (\exp x) \cdot J$. Since $A \subseteq J$ and $\exp_{\mathcal{FG}(B)}$ is a morphism of group germs, the map f is correctly defined and is a morphism of group germs; diagrammatic search shows that it is continuous. The canonical group monomorphism $\psi: G \rightarrow H$ is continuous as well, and $\psi \circ \phi = \exp_H \circ h$. Since for every $x \in \text{Lie}(H)$ one surely has $\exp_H(\lambda x) \neq e_H$ for an appropriate $\lambda \in \mathbb{R}$, the triple (\mathfrak{g}, G, f) is under conditions of Lemma 4.2. \square

Corollary 4.3. *Every continuous monomorphism from a Banach-Lie algebra with finite-dimensional center to the Lie algebra of a regular Lie group is tangent to an appropriate Lie group morphism.*

Proof. Results from direct application of Theorems 4.2 and 2.1. \square

As a corollary, we can now prove a restricted version of the Lie-Palais theorem by methods of regular Lie group theory.

Theorem 4.4 (Lie-Palais). *Let M be a compact closed manifold and let \mathfrak{g} be a finite-dimensional Lie algebra. Then every effective infinitesimal action of \mathfrak{g} on M is derived from a smooth global action of a finite-dimensional Lie group on M .*

Proof. The effective infinitesimal action of \mathfrak{g} upon M can be viewed as a Lie algebra monomorphism from \mathfrak{g} to the algebra $\text{vect } M$ of smooth vector fields on M endowed with the C^∞ -topology. Clearly, the conditions of the Main Theorem 4.2 and Corollary 4.3 are satisfied, and the resulting Lie group morphism from a Lie group G attached to \mathfrak{g} to the diffeomorphism group $\text{diff } M$ (which is a regular Lie group assigned to $\text{vect } M$, [KYMO, M]) determines a desired global smooth action. \square

Concluding Remarks 4.5. 1. We do not know whether the condition upon the center being finite-dimensional can be relaxed in the Main Theorem 4.2. However, our Example in [Pe2] can be remade easily to show that the Lemma 4.1 is no longer valid for Banach-Lie algebras with infinite-dimensional center.

2. Does not the present result shed a ray of hope, wee as it is, on an (apparently, abandoned) search for an “easy” proof of the Lie-Cartan theorem itself? (See [S], [H.V.8], and [Pe, Pe2].)

3. Another reputed open problem, which might well fall within this circle of ideas, is that of existence of an effective transitive action of a (connected) infinite-dimensional Banach-Lie group on a compact manifold [O, OdIH].

Acknowledgments. It is appropriate to thank Joshua Leslie for a few encouraging and stimulating letters sent by him in 1987 to the author who then worked at a secluded Siberian university. A Victoria University 1992-93 Small Research Grant is acknowledged.

REFERENCES

- [H] K.H. Hofmann, *Introduction to the Theory of Compact Groups. Part I*, Tulane University, Department of Mathematics, 1968.
- [KYMO] O. Kobayashi, A. Yoshioka, Y. Maeda and H. Omori, *The theory of infinite-dimensional Lie groups and its applications*, Acta Appl. Math. **3** (1985), 71–106.
- [L] J. Leslie, *Some integrable subalgebras of the Lie algebras of infinite-dimensional Lie groups*, Trans. Amer. Math. Soc. **333** (1992), 423–443.
- [M] J. Milnor, *Remarks on infinite-dimensional Lie groups*, Relativité, groupes et topologie II. Les Houches, Session XL, Elsevier Sci. Publ., Amsterdam a.o., 1984, pp. 1007–1058.
- [O] H. Omori, *Infinite Dimensional Lie Transformation Groups (Lecture Notes in Mathematics 427)*, Springer-Verlag, Berlin a.o., 1974.
- [OdIH] H. Omori and P. de la Harpe, *About interactions between Banach-Lie groups and finite dimensional manifolds*, J. Math. Kyoto Univ. **12** (1972), 543–570.
- [Pa] R.S. Palais, *A global formulation of Lie theory of transformation groups*, Mem. Amer. Math. Soc. **22** (1957).
- [Pe1] V.G. Pestov, *Free Banach-Lie algebras, couniversal Banach-Lie groups, and more*, Pacific J. Math. **157** (1993), 137–144.
- [Pe2] V.G. Pestov, *Enlargeable Banach-Lie algebras and free topological groups*, Bull. Austral. Math. Soc. **48** (1993), 13–22.
- [Po] M.M. Postnikov, *Lie Groups and Lie Algebras. Lectures in Geometry. Semester V*, Mir, Moscow, 1986 (Translated from Russian by V. Shokurov).
- [S] J.-P. Serre, *Lie Algebras and Lie Groups (1964 Harvard Lectures)*, 2nd ed. (Lecture Notes in Mathematics **1500**), Springer-Verlag, Berlin a.o., 1992.
- [Ś1] S. Świerczkowski, *Embedding theorems for local analytic groups*, Acta Math. **114** (1965), 207–235.
- [Ś2] S. Świerczkowski, *The path-functor on Banach-Lie algebras*, Indag. Math. **33** (1971), 235–239.
- [Ś3] S. Świerczkowski, *Cohomology of group germs and Lie algebras*, Pacific J. Math. **39** (1971), 471–482.
- [vEK] W.T. van Est and T.J. Korthagen, *Non-enlargible Lie algebras*, Nederl. Akad. Wetensch. Proc. **A26** (1964), 15–31.
- [vEŚ] W.T. van Est and S. Świerczkowski, *The path functor and faithful representability of Banach Lie algebras*, J. Austral. Math. Soc. **16** (1973), 471–482.